

Clinical Tests Reference Ranges

Items		Reference ranges		Units	Test Equipment and Analysis Method		Mesurement Facility
		Male	Female				
Biochemistry							
Protein,Total		6.7-8.3		g/dL	Beckman Coulter, Inc. AU5800	Biuret method	
Albumin		3.8-5.2		g/dL		Improved BCP method	
Albumin/Globulin Ratio		1.1-2.0					
Asparate aminotransferase		10-40		U/L		JSCC standardised method	
Alanine aminotransferase		5-45		U/L		JSCC standardised method	
Lactate dehydrogenase		124-222		U/L		IFCC standardised method	
Bilirubin,Total		0.2-1.2		mg/dL		Enzymatic assay	
Bilirubin,Direct		0.0-0.2		mg/dL		Enzymatic assay	
Bilirubin,Indirect		0.2-1.0		mg/dL			
Alkaline phosphatase		38-113		U/L		IFCC standardised method	
γ-Glutamyltransferase		≤80	≤30	U/L		JSCC standardised method	
Leucine aminopeptidase		45-81	37-61	U/L		L-leusyl-p-nitroanilid method	
Cholinesterase		234-493	200-452	U/L		JSCC standardised method	
Creatine kinase		60-270	40-150	U/L		JSCC standardised method	
Amylase		40-122		U/L		Enzymatic assay	
Blood urea nitrogen		8.0-20.0		mg/dL		Urease-GIDH method	
Creatinine		0.61-1.04	0.47-0.79	mg/dL		Enzymatic assay	
Uric acid		3.8-7.0	2.5-7.0	mg/dL		Enzymatic assay	
Sodium		137-147		mEq/L		ISE(Ion Selective Electorode)	
Chloride		98-108		mEq/L		ISE(Ion Selective Electorode)	
Potassium		3.5-5.0		mEq/L		ISE(Ion Selective Electorode)	
Calcium		8.4-10.4		mg/dL		Arsenazo III method	
Inorganic Phosphate		2.5-4.5		mg/dL		Enzymatic assay	
Magnesium		1.9-2.5		mg/dL		Enzymatic assay	
Cholesterol,Total		120-219		mg/dL		Enzymatic assay	
Cholesterol,LDL		65-139		mg/dL		Enzymatic assay (Direct method)	
Cholesterol,HDL		40-85	40-95	mg/dL		Enzymatic assay (Direct method)	
Triglyceride(s)		30-149		mg/dL		Enzymatic assay(Free glycerol remove method)	
Glucose		70-109		mg/dL		Enzymatic assay	
C-Reactive protein		≤0.30		mg/dL		Latex agglutination nephelometry	
HCO3-		22.6-31.4		mmol/L		Enzymatic assay	
Hematology							
White blood cell count		3300- 9000		/ μ L	Sysmex Corporation XN-1000	Flow Cytometry method	SUMIDA HOSPITAL
Red blood cell count		430- 570	380- 500	× 10 ⁴ / μ L		DC detection method	
Hemoglobin		13.5-17.5	11.5-15.0	g/dL		SLS-Hb method	
Hematocrit		39.7- 52.4	34.8- 45.0	%		DC detection method	
Mean corpuscular volume		85-102		fL			
Mean corpuscular hemoglobin		28.0- 34.0		pg			
Mean corpuscular hemoglobin concentration		30.2-35.1		%			
Platelet count		14.0-34.0		× 10 ⁴ / μ L		DC detection method	
Reticulocytes		4-19		%o		Flow Cytometry method	
Differential white blood cell count	Neutrophils	40.0-75.0		%		Flow Cytometry method	
	Lymphocytes	18.0-49.0		%			
	Monocytes	2.0-10.0		%			
	Eosinophils	0.0-8.0		%			
	Basophils	0.0-2.0		%			
	Neutrophils absolute count			× 10 ² / μ L			
	Lymphocytes absolute count			× 10 ² / μ L			
	Monocytes absolute count			× 10 ² / μ L			
Eosinophils absolute count			× 10 ² / μ L				
Basophils absolute count			× 10 ² / μ L				
Reticulocytes absolute count				× 10 ⁴ / μ L			
Urinalysis							
Color					Eye sight		Siemens Healthcare Diagnostics Corporation Clinitek Advantus
Specific gravity		1.006-1.030			Refractometer method		
pH		5.0-7.5			Testing paper		
Glucose		(-)					
Protein		(-)					
Occult blood		(-)					
Ketone bodies		(-)					
Bilirubin		(-)					
Urobilinogen		(+-)					
Nitrite		(-)					
Leukocytes		(-)					
Turbidity					Eye sight		
Sedimentary test,Erythrocytes				/HPF	Microscopic		
Sedimentary test,Leukocytes				/HPF			
Sedimentary test, Squamous epithelial cells				/HPF			
Drug testing,Urine		(-)			Gold colloids particle immunoassay		
Methadone test		(-)			Immuno-chromatography		
Pregnancy test					Immuno-chromatography		
Cotinine test		(-)			Immuno-chromatography		
Coaglation tests							
Prothrombin time;seconds		9.4-12.5		sec	IL JAPAN Corporation ACL TOP300	Coagulometricor mesurements	
Prothrombin time;Activity		70-100		%			
Prothrombin time;INR		0.85-1.15					
Activated partial thromboplastin time		25.0-36.0		sec			
Fibrinogen		155-415		mg/dL		clotting method adding thrombin	

Clinical Tests Reference Ranges

Items		Reference ranges		Units	Test Equipment and Analysis Method		Mesurement Facility	
		Male	Female					
Immunology								
Syphilis test,RPR		(-)			Beckman Coulter, Inc.	Latex agglutination nephelometry	SUMIDA HOSPITAL	
Syphilis test,TP antibody		(-)			AU5800	Latex agglutination nephelometry		
HBs antigen		(-)			Abbott Japan Corporation ARCHITECT2000	CLIA		
HCV antibody,2nd generation		(-)				CLIA		
HIV antigen and antibody		(-)				CLIA		
HBs antibody		(-)				CLIA		
HBc antibody		(-)				CLIA		
Others								
Breath alcohol test		0.00		mg/L	Semiconductor sensor			
Occult blood,Fecal		(-)			Immuno-chromatography			
SARS-CoV2		Negative			Tosoh Corporation TRCReady-80	TRC		
SARS-CoV-2 Agtest		(-)			Immuno-chromatography			
eGFR	Male:			mL/min/1.73m ²	194*Cre ^{-1.094} *age ^{-0.287}		SUMIDA HOSPITAL	
	Female:				194*Cre ^{-1.094} *age ^{-0.287} *0.739			
eGFR(EPI2021-J)	Male:			mL/min/1.73m ²	142*(Cre/0.9) ^{-0.302} *0.9938 ^{age} *0.813 (Cr<=0.9)			
					142*(Cre/0.9) ^{-1.2} *0.9938 ^{age} *0.813 (Cr>0.9)			
	Female:			mL/min/1.73m ²	142*(Cre/0.7) ^{-0.241} *0.9938 ^{age} *1.012*0.813(Cr<=0.7)			
					142*(Cre/0.7) ^{-1.2} *0.9938 ^{age} *1.012*0.813 (Cr>0.7)			
eGFR(EPI2021)	Male:			mL/min/1.73m ²	142*(Cre/0.9) ^{-0.302} *0.9938 ^{age} (Cr<=0.9)			
					142*(Cre/0.9) ^{-1.2} *0.9938 ^{age} (Cr>0.9)			
	Female:			mL/min/1.73m ²	142*(Cre/0.7) ^{-0.241} *0.9938 ^{age} *1.012 (Cr<=0.7)			
					142*(Cre/0.7) ^{-1.2} *0.9938 ^{age} *1.012 (Cr>0.7)			
GFR(MDRDJ)	Male:			mL/min/1.73m ²	175*Cre ^{-1.154} *age ^{-0.203} *0.808			
	Female:				175*Cre ^{-1.154} *age ^{-0.203} *0.742*0.808			
GFR(MDRD)	Male:			mL/min/1.73m ²	175*Cre ^{-1.154} *age ^{-0.203}			
	Female:				175*Cre ^{-1.154} *age ^{-0.203} *0.742			
CCR (Cockcroft-Gault)	Male:			mL/min	(140-age)*weight/(72*Cre)			
	Female:				(140-age)*weight/(72*Cre)*0.85			
Ca(Alb/adjust) Payne's formula				mg/dL	Ca+(4-ALB)			
Ca(Alb/adjust) KDOQI-2 formula				mg/dL	Ca+0.8×(4.0-ALB)			

Effective Date 01 Dec. 2025

SOUSEIKAI SUMIDA HOSPITAL